

FunSP, week 01

Ex 1 notes:

$$x[n] = u[n-20] - u[n-30]$$

unit step

$u[n]$




shifted unit step


$u[n-20]$

$$= 0$$

$$\therefore n=20$$




$u[n-30]$



$x[n]$




in Matlab:



NOTE: indexing in Matlab starts at 1 (not zero...)

Ex 3

$$h[n] = 2^{-n} u[n]$$



$$x[n] = u[n] - u[n-10]$$

$$y[n] = h[n] * x[n] = \sum_{k=-\infty}^{+\infty} h[k] x[n-k]$$


we will be using this alternative

$$= x[n] * h[n] = \sum_{k=-\infty}^{+\infty} x[k] h[n-k]$$

repeat the exercise at home using this other alternative...

First, we rename the independent variable as k :

$$h[k] = 2^{-k} u[k]$$



Second, we should understand what $x[n-k]$ is considering that k represents independent variable, and n represents a parameter because :

$$y[n] = \sum_k h[k] x[n-k]$$

↑ independent variable
↑ parameter
↑ independent variable




We understand that by noting that :

$$x[n-k] = x[-(k-n)]$$


which means that first we have $x[k]$, then $x[k-n]$ and, finally, $x[-(k-n)]$; in this process, we also need to understand what $x[-k]$ is with respect to $x[k]$

graphically we have:

(2)



which is the "mirror" version of $x[k]$ with respect to $k=0$, in the same way, $x[-(k-m)]$ is the "mirror" version of $x[k-m]$ with respect to $k-m=0$, i.e. with respect to $k=m$, thus, we have




this means that when the m parameter increases, the whole signal moves to the right and, when the m parameter decreases, the whole signal moves to the left

Now, we just need to find, for each value of the m parameter, the result of the element-wise product between the discrete-time sequences $h[k]$ and $x[m-k]$, and add all those products

we should address this by identifying intervals of values of m where the solution is given

by the same analytical expression.

For example, if $m < 0$ then we have




which makes it clear that the element-wise product between the two sequences is always zero, and, thus, the first interval of solutions is:

$m < 0$ $y[m] = 0$

Next, when $m \geq 0$, then there is an "overlap" between sequences $h[k]$ and $x[m-k]$, however, this overlap is either partial (when $m-9 < 0$) or total (when $m-9 \geq 0$), which gives rise to two new intervals:

$0 \leq m \leq 9$
could be < 9



In this case, we have:

$$y[m] = \sum_{k=0}^m h[k] x[m-k] = \sum_{k=0}^m 2^{-k} = \frac{1-2^{-(m+1)}}{1-\frac{1}{2}}$$

$$= 2(1-2^{-(m+1)}) = 2-2^{-m} = \frac{2^{m+1}-1}{2^m}$$


In this process, we should recall (and always keep in mind) the closed-form expression that gives the sum of terms of a geometric series:

$$\sum_{k=a}^b r^k = \frac{r^a - r^{b+1}}{1 - r}$$

Similarly, when

$$m \geq 9$$

$$y[m] = \sum_{k=m-9}^m h[k] r[m-k] \quad \text{because}$$



and, thus,

$$y[m] = \sum_{k=m-9}^m 2^{-k} = \frac{2^{-(m-9)} - 2^{-(m+1)}}{1 - \frac{1}{2}} = 2 \left(2^{-m+9} - 2^{-m-1} \right)$$

$$= \frac{2^{10} - 1}{2^{+m}}$$

An illustration of this three-interval solution is:

